

---

# TREATED SEED

This seed has been treated with **Axcess™ Insecticide Seed Treatment** containing **imidacloprid** at rates specified by the manufacturer.

## CAUTION

**SEED COMMERCIALY TREATED WITH AXCESS MUST BE  
LABELED IN COMPLIANCE WITH THE FEDERAL SEED ACT.**

### KEEP OUT OF REACH OF CHILDREN

**DO NOT** use treated seed for feed, food, or oil processing. Rapeseed greens and seed grown or harvested from **Axcess**-treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from **Axcess**-treated seed is only for industrial use and cannot be used for edible oil or any other human/feed consumption. Store away from feeds and other foodstuffs. Dispose of seed packaging in accordance with local requirements. Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow.

**DO NOT** induce vomiting unless told to do so by a poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

*(Continued on rear of tag.)*

NVA 2009-05-317-0154

2921467

---

BASF Corporation  
26 Davis Drive  
Research Triangle Park  
NC 27709

 **BASF**  
The Chemical Company

(Continued from front of tag.)

**ROTATIONAL PLANT-BACK INTERVALS\***

ROTATIONAL PLANT-BACK INTERVALS*		
Immediate Plant-Back	30-Day Plant-Back	
Artichoke	Mustard Seed	Cereals, including: buckwheat rice
Barley	Oats	
Brassica (cole) leafy veg.	Okra	
Borage	Pepinos	
Canola	Pepper	
Cilantro	Popcorn	
Corn, Field	Potato	
Corn, Sweet	Rapeseed	
Cotton	Rye	
Cranberry	Safflower	
Crambe	Sorghum	
Cucurbits	Soybean	
Eggplant	Strawberry	
Flax	Sugar beet	
Ground cherry	Sunflower	
Leaf petiole veg.	Tomatillo	
Leafy veg.	Tomato	
Legume veg. (succulent or dried, including soybean)	Triticale	
Millet	Veg. Root and Tuber	
	Watercress	
	Wheat	

\*Cover crops for soil building or erosion control may be planted any time, but **DO NOT** graze or harvest for food or feed.

\*For all other crops not listed on an Imidacloprid label, or for crops for which no tolerance for the active ingredient has been established, a 12-month plant-back interval is required.

**Applicators and other handlers must wear:** Long-sleeved shirt and long pants, socks and shoes and chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton). Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash hands and exposed skin thoroughly after handling the concentrate and after application.

**Use Restrictions:**

Rape greens grown and harvested from **Axcess**-treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from **Axcess**-treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.

**Wheat, Barley, Oats, Rye, Triticale, Sorghum, Millet: DO NOT** graze or feed livestock on treated areas for 45 days after planting.

**Cotton (delinted seed only):** Regardless of the type of application (seed treatment, soil or foliar) **DO NOT** apply more than a total of 0.5 lb. of imidacloprid per acre per cropping cycle.

**Seed-and-pod vegetable seed treated in California must be destined for planting in states other than California and is not to be planted in California.**